



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



Publication number: **0 646 644 A2**

12

## EUROPEAN PATENT APPLICATION

21 Application number: **94114325.7**

22 Date of filing: **12.09.94**

83 Declaration under Rule 28(4) EPC (expert solution)

51 Int. Cl.<sup>6</sup>: **C12N 15/62, C07K 14/28,  
C12N 9/12, C12N 1/21,  
C12N 5/10, G01N 33/68**

30 Priority: **13.09.93 US 121201**

43 Date of publication of application:  
**05.04.95 Bulletin 95/14**

84 Designated Contracting States:  
**AT BE CH DE DK ES FR GB GR IE IT LI LU MC  
NL PT SE**

71 Applicant: **BRISTOL-MYERS SQUIBB  
COMPANY  
P.O. Box 4000  
Princeton, NJ 08543-4000 (US)**

72 Inventor: **Menzel, Rolf  
1 Whitney Place  
Princeton Junction, NJ (US)  
Inventor: Taylor, Scott T.  
286, Wheatsheaf Lane  
Langhorne, PA (US)**

74 Representative: **VOSSIUS & PARTNER  
Postfach 86 07 67  
D-81634 München (DE)**

54 **Periplasmic membrane-bound system for detecting protein-protein interactions.**

57 The present invention relates to:

- (1) a fusion protein having a dimerizing domain with or without a ligand-binding region and toxR DNA-binding and hydrophobic transmembrane regions;
- (2) host cells comprising the fusion protein and a nucleic acid molecule having a reporter gene operatively linked to the ctx operon, wherein dimerization (ligand-dependent or -independent) is signaled by expression of the reporter gene;
- (3) a nucleic acid molecule coding for the fusion protein;
- (4) an expression vector comprising a coding region for the fusion protein;
- (5) a process for detecting dimer formation (ligand dependent or ligand independent) of the fusion protein, which comprises treating a culture of the host cells with a ligand, ligand mimetic, or dimerization inhibitor, and screening for expression of the reporter gene.

The present invention can be used to generate a signal from a variety of ligand-binding domains, allowing ligand binding to be indicated by a simple colorimetric test or antibiotic resistance. The fusion proteins could include therapeutically relevant domains, so that biologically pertinent interactions can be indicated by a readily measurable signal.

EP 0 646 644 A2

This invention relates to assays for protein-protein interactions, fusion proteins, and host cells modified to comprise such proteins.

Information about the extracellular environment is communicated to the inside of a cell by so-called signal transduction systems. Schlessinger, J., TIBS 13 443-7 (1988); Stock, J.B., et al., Microbiol. Rev. 53: 450-490 (1989). Several classes of signal transducers are able to "transmit" a signal without internalization of the primary signal. Such signal transmission may result from binding of a ligand to a membrane-bound receptor, inducing a conformational change in the receptor.

One such signal transducer is the toxR-ctx system in Vibrio cholerae. Ctx is the gene for cholera toxin. ToxR protein is a transmembrane protein with a DNA-binding domain at its amino terminus, a strongly hydrophobic transmembrane domain, and an environmentally regulated, dimer-forming domain at its carboxyl terminus. Miller, V. L., et al., Cell 48: 271-279 (1984). According to current beliefs, the hydrophobic transmembrane region directs the carboxyl terminus to the periplasm while the amino terminus remains in the cytoplasm. A change in osmolarity or temperature in the periplasm is believed to effect the capacity of the toxR protein to form dimers with its carboxyl terminus. The carboxyl dimers in turn modulate the ability of the amino terminus of toxR to bind to repetitive sequences in the ctx promoter and, thereby, modulate the ability of RNA polymerase to initiate transcription at the ctx promoter. Id.

To test the hypothesis that toxR protein is a membrane protein, Miller et al. fused toxR protein with alkaline phosphatase (phoA). Cell 48: 271-279 (1984). The phoA enzyme naturally exists as a dimer in the periplasm and the phoA-toxR fusion protein was constitutively functional with respect to both phosphatase activity and toxR activation of ctx. This result demonstrated that toxR was able to direct alkaline phosphatase to the periplasm where it was active, and that the dimerization capacity of alkaline phosphatase could direct the toxR protein to activate ctx transcription.

In a similar fusion experiment, Mekalanos et al. fused toxR to a phoA mutant that requires  $Zn^{2+}$  for dimerization and activity (personal communication). This fusion protein's phosphatase activity was dependent on  $Zn^{2+}$ , as expected, but so also was ctx activation. This result is consistent with the signal transduction model for toxR action.

Chimeric signal transduction systems have been constructed in prokaryotic cells with different bacterial systems (Utsumi, J., et al., Science 245: 1246-1249), in tissue culture cells with different eukaryotic hormone receptors (Riedel, H., et al., EMBO J. 8: 2943-2954 (1989)), and in tissue culture cells with a bacterial aspartate binding domain and the insulin receptor (Moe, G.E., et al., Proc. Natl. Acad. Sci. 86: 5683-7 (1989)). To date, however, no such system has employed the toxR receptor for V. cholerae.

The present invention relates to host cells comprising (a) a transmembrane fusion protein having (i) a periplasmic dimerization domain (e.g., a ligand-binding domain) and (ii) a toxR region having a cytoplasmic toxR DNA-binding region and a hydrophobic toxR transmembrane region, and (b) a nucleic acid molecule having a reporter gene operatively linked to the ctx operon, wherein dimer formation, which may or may not involve ligand binding, is signaled by expression of the reporter gene.

The present invention also relates to the aforementioned fusion protein, a corresponding nucleic acid molecule, and a corresponding expression vector having an inducible promoter. Further, the invention includes a process for detecting dimerization (e.g., as a result of binding of a ligand) of a membrane-bound protein, which comprises treating a culture of the host cells of this invention with a ligand and screening for expression of the reporter gene.

The present invention can be used to generate a signal from a variety of ligand-binding domains. Presence of a chosen ligand can be indicated by antibiotic resistance or a simple colorimetric test for production of an enzyme. The periplasmic regions of the transmembrane protein could include therapeutically relevant domains, so that biologically pertinent interactions can be indicated by a readily measurable signal. Such transmembrane fusion proteins should also contribute to our understanding of dimerization domains, ligand-binding domains, protein membrane-spanning sequences, and signal transduction.

Figure 1 is a schematic representation of the mechanism employed by the present invention. As shown in Figure 1, ligand-binding protein LBP is located within cell inner membrane IM and outer membrane OM. When the level expression of the toxR protein reaches a certain critical level, dimerization will ensue. Alternatively when a ligand L binds to the ligand binding domain of a toxR fusion protein, it may cause dimerization. In either instance, the dimerization induces a conformational change in the cytoplasmic domain of the wild type or toxR fusion protein which induces binding to the promoter region p of the reporter gene r. This DNA binding allows the RNA polymerase to transcribe the reporter gene, thus signalling dimerization which may result from either the spontaneous or ligand induced capacity of the periplasmic domain to associate.

Figure 2 shows control and fusion vectors. "LACp" denotes the *lac* promoter; "AmpR", the ampicillin resistance gene; "Spec", the spectinomycin resistance gene; "LacZ," the  $\beta$ -galactosidase gene; "CTXp", the cholera toxin promoter; "CAT", the chloramphenicol transferase gene. (A) pTOX3. This wild-type *ToxR* control vector includes a *toxR* PCR clone (SEQ. ID. NO. 1) from genomic *V. cholerae* in expression vector pSPORT1. The *toxR* gene is placed under the control of the *lac* promoter, making it inducible by treatment with IPTG. The desired transductants are selectable by treatment with ampicillin. The *ColEI* origin is compatible with the pSC101 origin. (B) pTOX5. This *toxR* fusion vector includes a truncated *toxR* PCR clone (SEQ. ID. NO. 2) from genomic *V. cholerae* in the vector pSPORT1. pTOX5 allows the in-frame cloning of foreign domains and may be used to generate families of fusions based on unidirectional deletion. The truncated pTOX5 does not activate the *ctx* promoter and any signals found in chimeric proteins result from the fused domain's capacity to dimerize. (C) pCTX3. This reporter gene vector includes the cholera toxin promoter, *ctx* (SEQ. ID. NO. 3), PCR-amplified from p3083 (Infection and Immunology, 58, pages 4142-4144, 1990) and cloned in a novel promoterless *lacZ* vector. This construct expresses  $\beta$ -galactosidase as a function of the dimerization of *ToxR* or *ToxR* fusion proteins. The pSC101 origin is compatible with the *ColEI* origin. The desired transformants may be selected by treatment with spectinomycin. (D) pCTX4. This vector includes a promoterless CAT gene from pCaMVCN inserted into the *LacZ* of pCTX3. Like the pCTX3 vector from which it is derived, pCTX4 also provides spectinomycin selection and expresses chloramphenicol acetyltransferase as a function of dimerization of *toxR* or *toxR* fusion proteins.

Figure 3 shows the results for the wild-type *toxR* gene, the truncated *toxR* from pTOX5, and a *toxR* in-frame gene fusion. The fusion in the example is to the leucine zipper dimerization domain of the yeast GCN4 protein, a well-studied dimerization domain representing an important control. (A) Co-transfection with *lacZ* reporter gene. For plasmids pTOX3, pTOX5, and pTOX5GCN4, this figure shows the amount of  $\beta$ -galactosidase produced (measured in Miller units) in the presence of zero (first bar), 50 (second bar), and 200 (third bar)  $\mu$ M IPTG. These results show that  $\beta$ -galactosidase expression is dependent on the induction of the wild type *toxR* gene of pTOX3 by IPTG, and demonstrate that the truncated *toxR* product in pTOX5 fails to activate expression even following induction. The results noted for pTOX5-GCN4 demonstrate that a foreign dimerization domain can restore inducible activation. (B) Co-transfection with CAT reporter gene. The results given in part B show the activation of the chloramphenicol resistance gene from pCTX4 as expected for the strains expressing the wild-type *toxR* and chimeric *toxR*-GCN4. The amount of chloramphenicol used is noted at the top of each column in  $\mu$ g/mL. "J4" denotes the strain JM101 as transfected with pCTX4. "R" signifies chloramphenicol resistance; "S", sensitivity. Cells were grown in the presence of IPTG as noted in the materials and methods; when IPTG is omitted, chloramphenicol resistance is not observed..

Figure 4 shows a method for generating unidirectional deletions, as applied to a *toxR*-*trkC* fusion. *TrkC* is a member of the tyrosine kinase class of receptors and interacts with the nerve growth factor NT3 (Cell, 66, pages 967-979, 1991). A full length *trkC* cDNA clone was placed downstream of the truncated *toxR* gene in the fusion vector pTOX5 at the *NotI* cleavage site. Following digestion with *SfiI* and *NotI*, this vector was treated with exonuclease III (*ExoIII*), which provides 5'-to-3' directional deletions starting at the *SfiI* cleavage site. *ExoIII*-treated plasmids were subsequently digested with mung-bean nuclease. Upon ligation, this treatment results in a family of fusions between *toxR* and *trkC* segments of varying length.

Figure 5 shows the results for the *toxR*-*trkC* fusions prepared by the procedure of Figure 4. (A).  $\beta$ -galactosidase assay. The *toxR* activity of various fusions was measured by  $\beta$ -galactosidase expression. Assays were performed in rich media (column LB) and minimal media (column min.) both in the presence of (100  $\mu$ M) and absence of IPTG. (B) Sequencing of *trkC* portion. The table shown in part B reports the size (in base pairs) and the boundaries of the sequence deleted in the fusion as determined by comparing the sequence of the deletion mutant with that of the known *trkC* and *toxR* sequences. (C). Map of *toxR*-*trkC* fusions. The data in the table in part B is expressed graphically with the hollow boxes indicating the base pairs deleted in the indicated fusion.

Figure 6 shows the results obtained with fusions of HSV ICP35 and HIV integrase to the *toxR* deletion of pTOX5. (A, B)  $\beta$ -galactosidase activity induced by *toxR* fusions. The results for a six-hour induction in JM101 background (A) and for a four-hour induction in DH5 $\alpha$  background (B) are shown. The y-axis shows  $\beta$ -galactosidase activity in Miller units; the x-axis, IPTG concentration in micromoles. These results confirm *ctx* activation by the *toxR*-viral fusion proteins and imply viral domain driven dimerization. (C) Chloramphenicol resistance induced by *ToxR* fusions. As in Figure 3, dimerization is demonstrated by resistance to chloramphenicol, reported in  $\mu$ g/mL. "D4" refers to strain DH5 $\alpha$  as a host strain and the remaining terms are as defined in Figure 3.

ToxR encodes a transmembrane protein possessing a DNA binding domain at its amino terminus. This intercellular domain has the capacity to activate expression from the *ctx* operon. Transactivation depends on the dimerization of *toxR* protein, which is directed by the protein's extracellular carboxyl domain. The *toxR-ctx* system can take the primary signal of protein-protein association and transduce it into the secondary message of gene expression. The natural membrane localization of the *toxR* protein, which is directed by its transmembrane segment, make it an attractive host for formation of chimera involving membrane proteins. This membrane confinement also has the intriguing feature of restricting protein diffusion to a 2-dimensional lipid plane. Placement of the dimerization domain in the extracellular periplasmic space will render it accessible to exogenous agents that might affect the dimer equilibrium.

#### Definition of terms

The following definitions apply to the terms as used throughout this specification, unless otherwise limited in specific instances.

The terms "dimerization domain" and "region capable of forming a dimer" refer to a polypeptide sequence capable of forming a self-dimer, either spontaneously or as a result of ligand binding and the like. Exemplary dimerization domains may be derived from *trkC*, GCN4, the HSV scaffold protein ICP35, HIV integrase, and the like.

The term "reporter gene" refers to any gene whose expression provides a measurable signal. Exemplary reporter genes include the genes for  $\beta$ -galactosidase and chloramphenicol transferase. Various other reporter genes are well known by those having ordinary skill in the art.

#### Process of preparation

##### Gene constructs

The nucleic acids used in the present invention may be prepared by recombinant nucleic acid methods. See, for example, the recombinant DNA methods of Nelles et al., *J. Biol. Chem.*, 262, 10855 (1987). Exemplary strains comprising such constructs are RFM2016 *supE thi Δ(pro-lac) /F' (traD36 proAB<sup>+</sup> lacI<sup>Q</sup> lacZΔM15) / pTOX5 / pCTX4* (ATCC Accession No. 69,403) and RFM2063 *supE thi Δ(pro-lac) degP::Tn5/F' (traD36 proAB<sup>+</sup> lacI<sup>Q</sup> lacZΔM15) / pTOX5trkC12 / pCTX3* (ATCC Accession No. 69,404). ("ATCC" refers to the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland 20852-1776.

The DNA sequences may be derived from a variety of sources, including genomic DNA, subgenomic DNA, cDNA, synthetic DNA, and combinations thereof. Genomic and cDNA may be obtained in a number of ways. Cells coding for the desired sequence may be isolated, the genomic DNA fragmented (e.g., by treatment with one or more restriction endonucleases), and the resulting fragments cloned, identified with a probe complementary to the desired sequence, and screened for the presence of a sequence coding for the desired activity.

For cDNA, the cDNA may be cloned and the resulting clone screened with a probe for cDNA coding for the desired region. Upon isolation of the desired clone, the cDNA may be manipulated in substantially the same manner as the genomic DNA.

To express the DNA sequences, transcriptional and translational signals recognized by an appropriate host are necessary. It is preferred that the *toxR* gene sequence be operatively linked to an inducible promoter (e.g., the *lac* promoter), providing a control for the system.

Alternatively, the promoter region from genomic DNA may be obtained in association with the DNA sequence for the *toxR* or other region for the fusion protein. To the extent that the host cells recognize the transcriptional regulatory and translational initiation signals associated with the *toxR* region, the 5' region adjacent to the coding sequence may be retained and employed for transcriptional and translational regulation. This region typically will include those sequences involved with initiation of transcription and translation, such as the TATA box, capping sequence, CAAT sequence, and the like. Typically, this region will be at least about 150 base pairs long, more typically about 200 bp, and rarely exceeding about 1 to 2 kb.

The non-coding 3' region may be retained, as well, especially for its transcriptional termination regulatory sequences, such as the stop signal and polyadenylated region. In addition, the non-coding 3' region may also contain an enhancer. Where the transcriptional termination signals are not satisfactorily functional in the host cell, then a functional 3' region from a different gene may be substituted. In this method, the choice of the substituted 3' region would depend upon the cell system chosen for expression.

A wide variety of transcriptional and translational regulatory sequences may be employed, depending upon the nature of the host. The transcriptional and translational regulatory sequences may be derived from viral sources (e.g., adenovirus, bovine papilloma virus, Simian virus, and the like) where the regulatory signals are derived from a gene that has a high level of expression in the host. Alternatively, promoters from mammalian expression products (e.g., actin, collagen, myosin, and the like) may be employed. Transcriptional initiation regulatory signals may be selected that allow for repression or activation, so that expression of the genes can be modulated. One such controllable modulation technique is the use of regulatory signals that are temperature-sensitive, so that expression can be repressed or initiated by changing the temperature. Another controllable modulation technique is the use of regulatory signals that are sensitive to certain chemicals.

To form the toxR or ctx chimeric gene constructs, DNA fragments may be ligated in accordance with conventional techniques known in the art. Such techniques include use of restriction enzymes to convert sticky-ended fragments to blunt ends (or vice-versa), polymerases and nucleotides to fill in sticky ends to form blunt ends, alkaline phosphatase to avoid undesired ligations, and ligases to join fragments.

The constructs for toxR and its fusion partner may be joined together to form a single DNA segment or may be maintained as separate segments by themselves or in conjunction with vectors. The constructs may be introduced into a cell by transformation in conjunction with a gene allowing for selection where the construct will become integrated into the host genome. Usually, the construct will be part of a vector having a replication system recognized by the host cell.

#### Expression vectors

Expression vehicles for production of the molecules of the invention include plasmids or other vectors. In general, such vectors contain control sequences that allow expression in various types of hosts, including but not limited to prokaryotes, yeasts, fungi, plants and higher eukaryotes. Suitable expression vectors containing the desired coding and control sequences may be constructed using standard recombinant DNA techniques known in the art, many of which are described in Sambrook, et al., Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1989).

An expression vector as contemplated by the present invention is at least capable of directing the replication of the reporter gene construct and the replication and expression of the transmembrane fusion protein construct. One class of vectors utilizes DNA elements that provide autonomously replicating extrachromosomal plasmids derived from animal viruses (e.g., bovine papilloma virus, polyomavirus, adenovirus, or SV40). A second class of vectors relies upon the integration of the desired gene sequences into the host cell chromosome.

Expression vectors useful in the present invention typically contain an origin of replication, a promoter located 5' to (i.e., upstream of) the DNA sequence to be expressed, and a transcription termination sequence. Suitable origins of replication include, for example, the ColE1, pSC101, SV40 and M13 origins of replication. Suitable termination sequences include, for example, the bovine growth hormone, SV40, lacZ and AcMNPV polyhedral polyadenylation signals. Suitable promoters include, for example, the cytomegalovirus promoter, the lacZ promoter, the gal 10 promoter and the AcMNPV polyhedral promoter. The promoter sequence may also be inducible, to allow modulation of expression (e.g., by the presence or absence of nutrients or other inducers in the growth medium). One example is the lac operon obtained from bacteriophage lambda plac5, which can be induced by IPTG.

The expression vectors may also include other regulatory sequences for optimal expression of the desired product. Such sequences include stability leader sequences, which provide for stability of the expression product; secretory leader sequences, which provide for secretion of the expression product; enhancers, which upregulate the expression of the DNA sequence; and restriction enzyme recognition sequences, which provide sites for cleavage by restriction endonucleases. All of these materials are known in the art and are commercially available. See, for example, Okayama, Mol. Cell. Biol., **3**, 280 (1983).

A suitable expression vector may also include marking sequences, which allow phenotypic selection of transformed host cells. Such a marker may provide prototrophy to an auxotrophic host, biocide resistance (e.g., antibiotic resistance) and the like. The selectable marker gene can either be directly linked to the DNA gene sequences to be expressed, or introduced into the same cell by co-transfection. Examples of selectable markers include neomycin, ampicillin, hygromycin resistance and the like.

The characteristics of the actual expression vector used must be compatible with the host cell that is to be employed. For a mammalian host, for example, the expression vector may contain promoters isolated from the genome of mammalian cells (e.g., mouse metallothionein promoter), or from viruses that grow in these cells (e.g., vaccinia virus 7.5 K promoter).

Suitable commercially available expression vectors into which the DNA sequences of the present invention may be inserted include pSPORT (which is preferred), pBluescriptII SK, the mammalian expression vectors pcDNAI or pcDNA/Neo, the baculovirus expression vector pBlueBac, the prokaryotic expression vector pcDNAII and the yeast expression vector pYes2, all of which may be obtained from Invitrogen Corp., San Diego, CA.

#### Host cells

The present invention additionally concerns host cells containing expression vectors that comprise DNA sequences for the toxR and ctx chimeric gene constructs.

Suitable host cells include both prokaryotic and eukaryotic cells. Suitable prokaryotic host cells include, for example, *E. coli* strains HB101, DH5a, XL1 Blue, Y1090 and JM101. Suitable eukaryotic host cells include, for example, *Spodoptera frugiperda* insect cells, COS-7 cells, human fibroblasts, and *Saccharomyces cerevisiae* cells.

Mammalian cells that may be useful as hosts include cells of fibroblast origin (e.g., VERO or CHO-K1) or lymphoid origin (e.g., SP2/0-AG14, or P3x63Sg8) or derivatives thereof. Preferred mammalian host cells include SP2/0 and J558L. Several cell lines secrete urokinase and may be used for transfection, such as cultured kidney carcinoma cells and 3T3 cells. Ferrivalo et al., *J. Cell Physiol.*, **121**, 363 (1984); Belin et al., *EMBO J.*, **3**, 190 (1984).

Another preferred host is yeast. Yeast provides substantial advantages in that it can also carry out post-translational peptide modification, including glycosylation. A number of recombinant DNA strategies exist that utilize strong promoter sequences and high copy number of plasmids that can be utilized for production of the desired proteins in yeast. Yeast recognizes leader sequences on cloned mammalian gene products and secretes peptides bearing leader sequences.

Immortalized cells, particularly myeloma or lymphoma cells, are also suitable host cells. These cells may be grown in an appropriate nutrient medium in culture flasks or injected into a syngenic host (e.g., mouse or rat) or an immunodeficient host or host site (e.g., nude mouse or hamster pouch). In particular, the cells may be introduced into the abdominal cavity for production of ascites fluid and harvesting of the chimeric molecule. Alternatively, the cells may be injected subcutaneously and the antibodies harvested from the blood of the host. The cells may be used in the same manner as the hybridoma cells. See Diamond et al., *N. Eng. J. Med.*, **304**, 1344 (1981); *Monoclonal Antibodies: Hybridomas-A New Dimension in Biologic Analysis* (Kennatt, et al., eds.) Plenum (1980).

Expression vectors may be introduced into host cells by various methods known in the art. For example, transfection of host cells with expression vectors can be carried out by the calcium phosphate precipitation method. However, other methods for introducing expression vectors into host cells, for example, electroporation, liposomal fusion, nuclear injection, and viral or phage infection can also be employed.

Host cells containing an expression vector may be identified by one or more of the following six general approaches: (a) DNA-DNA hybridization; (b) the presence or absence of marker gene functions; (c) assessing the level of transcription as measured by the production of mRNA transcripts encoding the gene constructs in the host cell; (d) detection of the gene product immunologically; (e) enzyme assay; and (f) PCR.

In the first approach, the presence of a DNA sequence coding for the gene constructs can be detected by DNA-DNA or RNA-DNA hybridization using probes complementary to the DNA sequence.

In the second approach, the recombinant expression vector host system can be identified and selected based upon the presence or absence of certain marker gene functions (e.g., thymidine kinase activity, resistance to antibiotics, etc.). A marker gene can be placed in the same plasmid as the fusion protein sequence or reporter construct under the regulation of the same or a different promoter or reporter. Expression of the marker gene indicates transfection of the vector having the DNA sequence for the fusion protein or reporter gene.

In the third approach, the production of mRNA transcripts encoding the fusion protein or reporter can be assessed by hybridization assays. For example, polyadenylated RNA can be isolated and analyzed by Northern blotting or a nuclease protection assay using a probe complementary to the RNA sequence. Alternatively, the total RNA of the host cell may be extracted and assayed for hybridization to such probes.

In the fourth approach, the expression of the fusion protein or reporter can be assessed immunologically, for example, by immunoblotting with antibody to the fusion protein or one of its regions (Western blotting). Alternatively, this technique could be carried out with the known epitope of the antibody variable region.

In the fifth approach, expression of the fusion protein or reporter can be measured by assaying for its activity (see below).

In the sixth approach, oligonucleotide primers homologous to sequences present in the expression system (i.e., expression vector sequences, fusion protein gene sequences, or reporter gene sequences) are used in a PCR to produce a DNA fragment of predicted length, indicating incorporation of the expression system in the host cell.

The expression vectors and DNA molecules of the present invention may also be sequenced. Various sequencing methods are known in the art. See, for example, the dideoxy chain termination method described in Sanger et al., *Proc. Natl. Acad. Sci. USA* 74, 5463-7 (1977), and the Maxam-Gilbert method described in *Proc. Natl. Acad. Sci. USA* 74, 560-4 (1977).

Once an expression vector has been introduced into an appropriate host cell, the host cell may be cultured under conditions permitting expression of large amounts of the desired protein.

The fusion protein may be isolated and purified in accordance with conventional conditions, such as extraction, precipitation, chromatography, affinity chromatography, electrophoresis, and the like. The preferred method is affinity chromatography with either the amino terminal heptapeptide of the fibrin  $\beta$  chain, which binds to the antifibrin site, or benzamidine, which binds to the plasminogen activator catalytic site.

It should, of course, be understood that not all expression vectors and DNA regulatory sequences will function equally well to express the DNA sequences of the present invention. Neither will all host cells function equally well with the same expression system. However, one of ordinary skill in the art may make a selection among expression vectors, DNA regulatory sequences, and host cells using the guidance provided herein without undue experimentation and without departing from the scope and spirit of the present invention.

#### Preferred Embodiments

Below are detailed descriptions of specific embodiments of the present invention. These embodiments are exemplary and serve to illustrate the broad applicability of the present invention.

The present inventors placed *V. cholerae*'s full length *toxR* gene behind the inducible *lac* promoter of the vector pSPORT. This provides a control to characterize the behavior of this system in *E. coli*. To exploit this system, the inventors cloned a truncated *toxR* gene immediately upstream of a multicloning site in the same parental vector (Figures 1 and 2).

#### TrkC Fusions

*ToxR-trkC* fusions were prepared as described in Figure 4. A large number of fusions with *trkC* segments of varying length were screened for *toxR* function ( $\beta$ -galactosidase activity at high IPTG in a host strain harboring the vector pCTX3). Clones with *toxR* activity were further characterized and the results are shown in Figure 5.

These fusions demonstrate that the foreign *trkC* sequences (without their own transmembrane segment) are able to restore varying levels of *toxR* function, which is fusion-dependent. Deletions that remove only small portions of *trkC* provide a fusion protein that demonstrates a *toxR* signal at higher levels of expression (high IPTG levels). In addition, a number of fusions that remove larger segments of *trkC* show enhanced dimer-forming ability, as demonstrated by *toxR* activity at low levels of expression (without IPTG.) In fact, many of the *trkC* fusions are more competent at *ctx* activation than is the wild type *toxR* protein with its own dimer forming domain. A possible model suggests that sequences in the wild-type *trkC* amino terminus obscure a very strong dimer forming capacity. In this model, NT3 binding in the wild type *trkC* protein might cause conformational changes that uncover this strong self-association domain.

The modified host cells allow a screen for *trkC* ligands with NT3-like activity, using chloramphenicol resistance or  $\beta$ -galactosidase as an indicator. Ligand activity will be signaled by the appearance of chloramphenicol resistance or  $\beta$ -galactosidase activity at a level of fusion expression below that which gives ligand-independent activity.

#### Other fusions

In addition to *trkC* and GCN4-directed *toxR* association, both the HSV scaffold protein ICP35 and the HIV integrase protein are competent to direct *toxR* function (see Figure 6). Other suitable periplasmic binding regions for this invention include the bacterial chemotaxis receptors tar (aspartate) and tsr (serine), the human insulin receptor and HIV protease. All such fusions will allow screens to identify inhibitors



specific to their respective fused dimerization domains.

## Materials and methods

### 5 Strains

Vibrio cholera strain 569B rpf, classical biotype, Inaba serotype (courtesy of H.Smith) (BMS culture collection number SC15307) was used for all V. cholera DNA isolations.

E. coli strain JM101 (Messing 1979) was used in all ctx transactivation experiments. E. coli strains DH5 $\alpha$  (Gibco-BRL) and JM101 were used for general plasmid maintenance and amplification.

### Media

Vibrio strains were grown in NB (nutrient broth) supplemented with 0.5% NaCl at 30 °C. E. coli strains were grown in LB (Luria-Bertani) broth at 37 °C. When selection of plasmids was required, ampicillin was used at 50  $\mu$ g/mL and spectinomycin was used at 50  $\mu$ g/mL. Chloramphenicol was used at varying concentrations as described in the text.

### Genetic techniques

Standard methods were used for the isolation and manipulation of plasmid DNA (Maniatis, T., Fritsch, E.F. & Sambrook, J. (1982) Molecular Cloning: A Laboratory Manual (Cold Spring Harbor Lab., Plainview, NY). Genomic Vibrio DNA was prepared as described in Clark, J.M. & Switzer, R.L. (1977) Experimental Biochemistry, p. 230 (W.H. Freeman Co. NY).

Unidirectional deletions were formed using ExoIII, mung bean nuclease, T4 DNA ligase from a unidirectional deletion kit (Stratagene) according to protocols supplied by the manufacturer. Briefly, the plasmid pTOX5TRKC was digested with SfiI and NotI restriction endonucleases prior to the addition of ExoIII. Aliquots were taken at one minute intervals for 5 minutes from the ExoIII mixture and the plasmid ends were flushed with mung bean nuclease and then ligated to create deletions into the trkC portion of the toxR-trkC chimera. These plasmids were transformed into JM101/pCTX3 (J3) and assayed for  $\beta$ -galactosidase as a measure of their ability to transactivate ctx-LACZ. Active fusions such as pTOX5TRKC  $\Delta$ 12 were then sequenced (B. Keinzle) on an ABI DNA sequencer to define the deletion endpoints and verify that an in-frame fusion was generated.

DNA was amplified by PCR using a Perkin-Elmer Cetus thermal DNA cycler 480 and GeneAmp DNA amplification kit. The reaction mix contained 1X Amplitaq Buffer with 1.5 mM magnesium chloride, 200  $\mu$ M each dNTP, 2.5 U Amplitaq polymerase, 1.0  $\mu$ M 5' and 3' primers and 10-100 ng template DNA in 100  $\mu$ L total volume. In general, the amplification consisted of one cycle of 5 minutes at 94 °C, 35 cycles consisting of one minute at 94 °C followed by 2 minutes at 52 °C then 3 minutes at 72 °C and completed with a final cycle of 8 minutes at 72 °C. PCR products are analyzed on 0.8% agarose gels and then digested with restriction endonucleases (at sites engineered into the primers) prior to ligation into the host vector.

Standard methods were used for SDS-PAGE and immunoblotting of toxR fusion proteins (Maniatis, T., Fritsch, E.F. & Sambrook, J. (1982) Molecular Cloning: A Laboratory Manual (Cold Spring Harbor Lab., Plainview, NY). Generally, whole cell protein or purified protein was loaded onto a 10% SDS Daiichi mini gel and electrophoresis was run at 40 mA constant current in 1X Laemmli buffer for one hour. Following electrophoresis, the proteins were transferred to a supported nitrocellulose membrane in a Hoeffer blotting tank for one to two hours at 375 mA constant current. Western analysis was conducted using a 1:1000 dilution of the primary antibody (rabbit anti-GSTTOX5 polyclonal sera) followed by a second incubation with a 1:3000 dilution of the secondary antibody (mouse or goat anti-rabbit monoclonal antibodies, BioRad) conjugated to alkaline phosphatase. The membrane was blocked with 5% powdered skim milk in Tris buffered saline (blotto) before the addition of the primary antibody and washed three times with blotto after the addition of each antibody. Antigenic proteins were visualized using the western blue stabilized substrate for alkaline phosphatase (Promega).

### 55 Plasmid Vectors

pCTX3 and pCTX4. The plasmid pCTX3 was constructed by digesting pMLB1109 with PstI and BalI, and isolating the appropriate 5 kb fragment. This fragment was then ligated to a 5.7 Kb PstI and SmaI



fragment of pMS421. The resulting plasmid, pCTX2, was partially digested with EcoRI to delete the 1.7 Kb lacI<sup>q</sup> region. The EcoRI ends from the partial were filled in with Klenow polymerase and dNTPs, and then ligated to yield the vector pSLAC2. The plasmid pCTX3 was constructed by cloning an EcoRI-BamHI flanked PCR fragment of the CTX promoter from p3083 (R. Finklestein) into EcoRI and BamHI digested pSLAC2 to yield the 9.1 Kb plasmid. The plasmid pCTX4 was generated by inserting the BamHI and ClaI digested fragment of pCaMVCN (Pharmacia) containing the CAT gene into BamHI and ClaI digested pCTX3.

pTOX3, pTOX4, and pTOX5. The plasmid pTOX3 was constructed by PCR amplifying the toxR coding region with a 5' PstI restriction site containing PCR primer and a 3' SfiI restriction site harboring PCR primer using genomic DNA prepared from the V. cholera strain 569B rpf obtained from H. Smith. The resulting PCR fragment was cloned into PstI- and SfiI-digested pSPORT1 (BRL).

The plasmid pTOX4 was constructed by PCR amplifying, from genomic V. cholera DNA, the toxR coding region truncated at base pair 989 of the toxR reading frame using a PCR primer with a 5' PstI restriction site and a second primer with both a 3' SfiI restriction sites and stop codons. The resulting PCR fragment was cloned into PstI- and SfiI-digested pSPORT1 (BRL).

The plasmid pTOX5 was constructed by PCR amplifying a truncated version of the toxR coding region at base pair 827 of the toxR reading frame in a manner strictly analogous to that used in the pTOX4 construct.

pTOX5ICP35. The plasmid pTOX5ICP35 was developed by digesting pT7ICP35K (I. Deckman) with KpnI and NcoI to yield a fragment containing the HSV ICP35 coding region. The NcoI site of this fragment was filled in with Taq polymerase and dNTPs. This modified insert was cloned into KpnI and StuI digested pTOX5 giving a fusion of the ICP35 reading frame aligned with that of the truncated toxR gene.

pTOX5TRKC+TM. The plasmid pTOX5TRKC+TM was constructed by generating a 1.3 kb PCR fragment of the trkC coding region from base pair position 32 to position 1391 of the trkC reading frame of pFL19 (M. Barbacid) containing 3' XbaI and 5' HindIII termini with stop codons derived from the PCR primers. This fragment was then cloned (not in-frame) into XbaI and HindIII digested pTOX5 to create a vector that could be used to make unidirectional deletions of trkC truncated just past the transmembrane domain fused to toxR.

pTOX5TRKC-TM. The plasmid pTOX5TRKC-TM was generated in the same fashion as pTOX5TRKC+TM, but the 1.3 kb PCR-generated (trkC position 32 to 1318) fragment did not contain the trkC transmembrane domain.

pTOX5TRKC-TM del 12. The plasmid pTOX5TRKC-TM  $\Delta$ 12 was created by treating with ExoIII the SfiI- and NotI-digested pTOX5TRKC-TM. ExoIII treatment was stopped at one-minute intervals, and mung bean nuclease was added to flush the ends prior to rejoining the plasmid deletions with T4 ligase. Deletion number 12 was sequenced and shown to be an in-frame deletion toxR-trkC segment of pTOX5TRKC-TM. The junction of the deletion contains the amino acid residues TG from the end of the truncated toxR gene with the intervening linker RSRSE joined to the initial methionine of the trkC sequence.

pTOX5HIVPROT. The plasmid pTOX5HIVPROT was devised from an SfiI-flanked 300-base pair PCR-amplified HIV Protease gene (P-F. Lin) cloned in-frame into SfiI-digested pTOX5.

pTOX5HIVINT. The plasmid pTOX5HIVINT was formulated by inserting an 896-base pair PCR product of the HIV integrase gene, with an SfiI site at the 5' terminus and XbaI site at the 3' terminus derived from the PCR primers (P-F. Lin), into SfiI and XbaI-digested pTOX5 to yield an in-frame fusion of the HIV integrase with toxR.

pTOX5GCN4. The plasmid pTOX5LINK was established to allow in-frame cloning of the leucine zippers of the yeast vesulatory protein GCN4 with SalI and BamHI termini. This vector was created by digesting pTOX5 with SfiI and BamHI, and cloning in two annealed oligonucleotides (sequences 5' to 3' GTCGACG and GATCCGTCGACCTC). The plasmid pTOX5GCN4 was originated by liberating a 129-bp fragment from SalI- and BamHI-digested pJH370 (J. Hu) and cloning this piece into SalI- and BamHI-digested pTOX5LINK.

pTOX5LP19. pTOX5LV19. pTOX5LY19. pTOX5LI19. pTOX5LN19. pTOX5LK19. The plasmids pTOX5LP19, pTOX5LV19, pTOX5LY19, pTOX5LI19, pTOX5LN19, pTOX5LK19 were engineered identically to pTOX5GCN4 but included the SalI-BamHI fragments from pJH524, pJH505, pJH506, pJH518, pJH521, pJH528 (J. Hu) respectively.

pGSTTOX5. The plasmid pGSTTOX5 was made by digesting pGEX-3X (Pharmacia) with SmaI and BamHI and cloning in PCR amplified truncated toxR from pTOX5 with compatible ends.

Assays

$\beta$ -galactosidase assays were performed essentially as described by Menzel (Analytical Biochemistry 181, p. 40-50 1989.)

5 Agar plate assays for ctx activation based on chloramphenicol resistance were performed with JM101 strains containing the plasmid pCTX4 and any of the following activating plasmids; pTOX3, pTOX5, pTOX5GCN4, pTOX5ICP35, or pTOX5HIVINT. Overnight cultures were grown in LB with 100  $\mu$ M IPTG. One to two microliters of the overnight culture was spotted on LB-agar plates with 100  $\mu$ M IPTG and the levels of chloramphenicol indicated in the text. ToxR activity was scored based on the ability to grow on  
10 increasing levels of the drug. In the absence of IPTG, all strains failed to show any chloramphenicol resistance.

The abbreviations used in this specification are defined as follows.

	AcMNPV	<u>Autographa californica</u> multiple nuclear polyhedrosis virus
	bp	base pairs
15	CAT	chloramphenicol transferase
	cDNA	complementary DNA
	DNA	deoxyribonucleic acid
	HIV	human immunodeficiency virus
	HSV	herpes simplex virus
20	IPTG	isopropylthiogalactoside
	kb, kbp	kilobase pairs
	NT	neurotrophin
	PCR	polymerase chain reaction
	phoA	alkaline phosphatase
25	RNA	ribonucleic acid

30

35

40

45

50

55

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- 5 (i) APPLICANT: Menzel. Rolf  
Taylor. Scott T.
- 10 (ii) TITLE OF INVENTION: PERIPLASMIC MEMBRANE-BOUND SYSTEM FOR  
DETECTING PROTEIN-PROTEIN INTERACTIONS
- (iii) NUMBER OF SEQUENCES: 3
- (iv) CORRESPONDENCE ADDRESS:  
15 (A) ADDRESSEE: Burton Rodney  
(B) STREET: P.O. Box 4000  
(C) CITY: Princeton  
(D) STATE: New Jersey  
(E) COUNTRY: U.S.A.  
(F) ZIP: 08543-4000
- 20 (v) COMPUTER READABLE FORM:  
(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- 25 (vi) CURRENT APPLICATION DATA:  
(A) APPLICATION NUMBER:  
(B) FILING DATE:  
(C) CLASSIFICATION:
- 30 (viii) ATTORNEY/AGENT INFORMATION:  
(A) NAME: Gaul, Timothy J.  
(B) REGISTRATION NUMBER: 33.111  
(C) REFERENCE/DOCKET NUMBER: DC28
- 35 (ix) TELECOMMUNICATION INFORMATION:  
(A) TELEPHONE: (609) 252-5901  
(B) TELEFAX: (609) 252-4526
- 40
- 45
- 50
- 55

EP 0 646 644 A2

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1105 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

	GTGAGTGAGT GAGTGTGGG ACAGGGACAT ACTGGGACAT TAGATGTTTCG GATTAGGACA	60
	CAACTCAAAA GAGATATCGA TGAGTCATAT TGGTACTAAA TTCATTCTTG CTGAAAAATT	120
15	TACCTTCGAT CCCCTAAGCA ATACTCTGAT TGACAAAGAA GATACTGAAG AGATCATTCG	180
	ATTAGGCAGC AACGAAAGCC GAATTCCTTG GCTGCTGGCC CAACGTCCAA ACGAGGTAAT	240
	TTCTCGCAAT GATTTGCATG ACTTTGTTTG GCGAGAGCAA GGTTTGAAG TCGATGATTC	300
20	CAGCTTAACC CAAGCCATTT CGACTCTGCG CAAAATGCTC AAAGATTCGA CAAAGTCCCC	360
	ACAATACGTC AAAACGGTTC CGAAGCGCGG TTACCAATTG ATCGCCCGAG TGGAAACGGT	420
	TGAAGAAGAG ATGGCTCGCG AAAACGAAGC TGCTCATGAC ATCTCTCAGC CAGAATCTGT	480
25	CAATGAATAC GCAGAATCAA GCAGTGTGCC TTCATCAGCC ACTGTAGTGA ACACACCGCA	540
	GCCAGCCAAT GTCGTGGCGA ATAAATCGGC TCCAAACTTG GGAATCGAC TGTTTATTCT	600
	GATAGCGGTC TTACTIONCCCC TCGCAGTATT ACTGCTCACT AACCCAAGCC AATCCAGCTT	660
30	TAAACCCCTA ACGGTTGTCG ATGGCGTAGC CGTCAATATG CCGAATAACC ACCCTGATCT	720
	TTCAAATTGG CTACCGTCAA TCGAAGTGTG CGTTAAAAAA TACAATGAAA AACATACTGG	780
	TGGACTCAAG CCGATAGAAG TGATTGCCAC TGGTGGACAA AATAACCACT TAACGCTGAA	840
35	TTACATTCAC AGCCCTGAAG TTTCAGGGGA AAACATAACC TTACGCATCG TTGCTAACCC	900
	TAACGATGCC ATCAAAGTGT GTGAGTAGGA TCTTGCTATG CAAAATAGAC ACATCGCCAT	960
	GGGTATTCTT CATAGGAAAA CTGAAGAGCA TCTGATCGAC TTCATATCA CAGTTCCCAC	1020
40	GCACAGCAAT GATCTGCTGA GCAAACGAT TCAATTTTTC AGCGACGGCT ACAGGGTTGT	1080
	ACCTTGCGGG CCATCTAGGC CTGCA	1105

EP 0 646 644 A2

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 843 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

TTCCTGCAGT	GAGTGAGTGA	GTGTTGGGAC	AGGGAGATAC	TGGGACATTA	GATGTTTCGGA	60
TTAGGACACA	ACTCAAAAGA	GATATCGATG	AGTCATATTG	GTACTAAATT	CATTCTTGCT	120
GAAAAATTTA	CCTTCGATCC	CCTAAGCAAT	ACTCTGATTG	ACAAAGAAGA	TAGTGAAGAG	180
ATCATTCGAT	TAGGCAGCAA	CGAAAGCCGA	ATTCTTTGGC	TGCTGGCCCA	ACGTCCAAAC	240
GAGGTAATTT	CTCGCAATGA	TTTGCATGAC	TTTGTTTGGC	GAGAGCAAGG	TTTTGAAGTC	300
GATGATTCCA	GCTTAACCCA	AGCCATTTTC	ACTCTGCGCA	AAATGCTCAA	AGATTTCGACA	360
AAGTCCCCAC	AATACGTCAA	AACGGTTCGC	AAGCGCGGTT	ACCAATTGAT	CGCCCGAGTG	420
GAAACGGTTG	AAGAAGAGAT	GGCTCGCGAA	AACGAAGCTG	CTCATGACAT	CTCTCAGCCA	480
GAATCTGTCA	ATGAATACGC	AGAATCAAGC	AGTGTGCCTT	CATCAGCCAC	TGTAGTGAAC	540
ACACCGCAGC	CAGCCAATGT	CGTGGCGAAT	AAATCGGCTC	CAAACTTGGG	GAATCGACTG	600
TTTATTCTGA	TAGCGGTCTT	ACTTCCCCTC	GCAGTATTAC	TGCTCACTAA	CCCAAGCCAA	660
TCCAGCTTTA	AACCCCTAAC	GGTGTGCGAT	GGCGTAGCCG	TCAATATGCC	GAATAACCAC	720
CCTGATCTTT	CAAATTGGCT	ACCGTCAATC	GAAGTGTGCG	TTAAAAAATA	CAATGAAAAA	780
CATACTGGTG	GACTCAAGCC	GATAGAAGTG	ATTGCCACTG	GTGGCCATCT	AGGCCTGCAG	840
GAA						843

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 145 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

AACAGAAAAT	GATAAAAAAG	GACTAATAGT	ATATTTTGAT	TTTTGATTTT	TGATTTTGA	60
TTTTTGATTT	TTGATTTTGG	ATTTTGTGAT	TCAAATAATA	CAAATTTATT	TACTTATTTA	120
ATTGTTTGA	TCAATTATTT	TTCTG				145

Claims

1. A host cell comprising
  - (a) a transmembrane fusion protein having
    - (i) a toxR region having a toxR DNA-binding region and a toxR hydrophobic transmembrane region, and
    - (ii) a periplasmic region capable of forming a dimer spontaneously or upon binding of a ligand; and
  - (b) a nucleic acid molecule having a reporter gene operatively linked to the ctx operon;
 wherein dimer formation is signaled by expression of the reporter gene.
2. A transmembrane fusion protein, comprising
  - (a) a toxR region having a toxR DNA-binding region and a toxR hydrophobic transmembrane region, and
  - (b) a periplasmic region capable of forming a dimer spontaneously or upon binding of a ligand.
3. A nucleic acid molecule coding for the transmembrane fusion protein of Claim 2.
4. An expression vector, comprising the nucleic acid of Claim 3.
5. The host cell of Claim 1, wherein the periplasmic region forms a dimer upon binding of a ligand.
6. A process for detecting binding of a ligand to a periplasmic region capable of forming a dimer upon binding of a ligand, which comprises:
  - (a) treating a culture of the host cells of Claim 5 with a ligand, and
  - (b) screening for expression of the reporter gene.
7. The host cell of Claim 1, wherein the toxR region has the sequence of SEQ. ID. NO. 1.
8. The chimeric transmembrane protein of Claim 2, wherein the toxR region has the sequence of SEQ. ID. NO. 1.
9. The nucleic acid of Claim 3, wherein the toxR region has the sequence of SEQ. ID. NO. 1.
10. The expression vector of Claim 4, wherein the toxR region has the sequence of SEQ. ID. NO. 1.
11. The process of Claim 5, wherein the toxR region has the sequence of SEQ. ID. NO. 1.
12. The nucleic acid of Claim 3, wherein the toxR region has the nucleic acid sequence of SEQ. ID. NO. 2.
13. The expression vector of Claim 4, wherein the toxR region has the nucleic acid sequence of SEQ. ID. NO. 2.
14. The process of Claim 6, wherein the toxR region has the nucleic acid sequence of SEQ. ID. NO. 2.
15. The host cell of Claim 1, designated ATCC 69,403 or ATCC 69,404.
16. The host cell of Claim 1, wherein the periplasmic region comprises the ligand-binding region of trkC.
17. The protein of Claim 2, wherein the periplasmic region comprises the ligand-binding region of trkC.
18. The nucleic acid of Claim 3, wherein the periplasmic region comprises the ligand-binding region of trkC.
19. The expression vector of Claim 4, wherein the periplasmic region comprises the ligand-binding region of trkC.
20. The process of Claim 6, wherein the periplasmic region comprises the ligand-binding region of trkC.






FIG. 2A




FIG. 2B




FIG. 2C






FIG. 2D



**FIG. 3A****FIG. 3B**

		[Chloramphenicol] μg/ml				
		0.00	6.25	12.00	25.00	50.00
Strain	J4/pTOX3	R	R	R	R	R
	J4/pTOX5	R	S	S	S	S
	J4/pTOX5GCN4	R	R	R	R	S

J4= JM101/pCTX4  
 R= Resistant  
 S= Sensitive





FIG. 5A

	LB-	LB+	MIN-	MIN+	
+TM,DEG+	trkC1	2087	1635	3519	2564
	trkC2	1118	1391	1577	1939
	trkC3	1454	4987	2277	8363
	trkC4	1515	4221	2162	6817
	trkC5	971	725	1501	1067
	trkC6	798	3238	1078	5223
	trkC7	891	881	1303	1292
-TM,DEG-	trkC8	2322	1380	2978	2537
	trkC9	777	643	1034	1078
	trkC10	3267	3154	3702	4688
	trkC11	1163	818	1592	1403
	trkC12	1015	3389	1315	7263
	trkC13	956	807	1509	1349
	trkC14	2872	2953	4199	4307
	trkC15	863	4065	1165	6804
	trkC16	602	1757	1068	4775
	trkC17	821	4304	1396	6771
	trkC18	982	2057	1413	3214
	trkC19	940	3261	1418	5283
	trkC20	1075	932	1774	1503
-TM,DEG+	trkC21	931	1310	1692	1939
	trkC22	2525	4366	3069	6165
	trkC23	2249	2445	2919	3819
	trkC24	850	2751	1181	3497
	trkC25	1391	1766	2409	3058
	trkC26	898	3494	1246	4326
	trkC27	651	3539	1216	5655
	trkC28	1156	3044	1659	3948
	trkC29	447	2222	579	2013
	trkC30	672	1277	1032	1435
Controls					
	tox3	554	2443	903	4253
	tox3	609	2888	895	4638
	tox3	524	2818	920	4144
	tox5	284	286	664	711
	tox5	365	240	661	683
	302	274	790	625	

FIG. 5B

trkC Deletion Number	Deletion Size (bp)	Deletion Endpoints in pTOXTRKC-tm
trkC 1	875	634 - 1509
trkC 3	905	634 - 1539
trkC 6	389	634 - 1023
trkC 10	683	634 - 1317
trkC 12	71	634 - 705
trkC 14	347	639 - 986
trkC 15	71	634 - 705
trkC 17	71	634 - 705
trkC 22	71	634 - 705
trkC 23	407	634 - 1041

FIG. 5C






FIG. 6A




FIG. 6B

Strain	[Chloramphenicol] $\mu\text{g/ml}$					
	0.00	2.50	5.00	10.00	20.00	40.00
	D4/pTOX3	R	R	R	R	R
	D4/pTOX5	R	R	S	S	S
	D4/pTOX5ICP35	R	R	R	S	S
	J4/pTOX3	R	R	R	R	R
	J4/pTOX5	R	R	S	S	S
	J4/pTOX5HIVINT	R	R	R	R	S

FIG. 6C

D4= DH5 $\alpha$   
 J4= JM101  
 R= Resistant  
 S= Sensitive





European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 94 11 4325

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
D,X	CELL, vol. 48, 30 January 1987, pages 271-279, XP002002026 MILLER, V.L. ET AL.: "Cholera toxin transcriptional activator toxR is a transmembrane DNA binding protein" page 272, right-hand column, third paragraph; Figure 2; page 274, left-hand column	1-6	C12N15/62 C07K14/28 C12N9/12 C12N1/21 C12N5/10 G01N33/68
Y	whole document	7-20	
X	CELL, vol. 64, 11 January 1991, pages 29-34, XP002002027 DIRITA, V.J. AND MEKALANOS, J.J.: "Periplasmic interaction between two membrane regulatory proteins, toxR and toxS, results in signal transduction and transcriptional activation" page 31, right-hand column, second paragraph	1-6	
Y	whole document	7-20	
E	EP-A-0 630 968 (BEHRINGWERKE) 28 December 1994 pages 5-6; Example III	1-6	
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 3 May 1996	Examiner Alt, G
<p><b>CATEGORY OF CITED DOCUMENTS</b></p> <p>X : particularly relevant if taken alone  Y : particularly relevant if combined with another document of the same category  A : technological background  O : non-written disclosure  P : intermediate document</p> <p>T : theory or principle underlying the invention  E : earlier patent document, but published on, or after the filing date  D : document cited in the application  L : document cited for other reasons  &amp; : member of the same patent family, corresponding document</p>			

EPO FORM 1501 03/92 (P0409)



European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 94 11 4325

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
P,X	BIOL.CHEM.HOPPE-SEYLER, vol. 375, January 1994, pages 61-70, XP002002028 KOLMAR, H. ET AL.: "Dimerization of Bence Jones proteins: Linking the rate of transcription from an Escherichia coli promoter to the association constant of REIV" page 66, right-hand column, line 11 - page 67, left-hand column, line 12; Figure 6 ---	1-6	
D,A	CELL, vol. 66, 6 September 1991, pages 967-979, XP002002029 LAMBALLE, F. ET AL.: "trkC, a new member of the trk family of tyrosine protein kinases, is a receptor for neurotrophin-3" see whole document -----	16-20	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
Place of search MUNICH		Date of completion of the search 3 May 1996	Examiner Alt, G
<p><b>CATEGORY OF CITED DOCUMENTS</b></p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons * : member of the same patent family, corresponding document</p>			

EPO FORM 1503 01.92 (P04/D1)